OXYMETOS 25 mg Generic Medicine (100 tabs). Real Oxymetholone



Product Name: OXYMETOS 25 mg
Category:Oral Steroids
Ingredient: Oxymetholone
Manufacturer: Pharmacom
Qty: 100 tabs
Buy online: https://t.co/Gmuee0I43C

Tablets. 12.5 mg; 25 mg; 32 mg; 50 mg; Tablet, chewable. 25 mg; Dosage Considerations - Should be Given as Follows: Motion Sickness. 12.5-50 mg orally given 1 hour before travel and then every 24 hours as needed; Children older than 12 years: 12.5-50 mg orally given 1 hour before travel and then daily as needed Find patient medical information for Maxzide-25mg oral on WebMD including its uses, side effects and safety, interactions, pictures, warnings and user ratings.



Side Effects. Hot flashes, hair loss, joint /bone/ muscle pain, tiredness, unusual sweating, nausea, diarrhea, dizziness, and trouble sleeping may occur. If any of these effects persist or worsen. Metoprolol 25 MG Tablet slows down the activity of your heart, allowing your heart to beat less forcefully. Thus, this medicine helps to prevent abnormally fast heartbeats. It also reduces the blood pressure in your body. Metoprolol 25 MG Tablet may cause some side effects like dizziness, headache, breathing difficulty, and low blood pressure.



Diclofenac sodium enteric-coated and delayed-release tablets: 25 mg orally 4 times a day. An additional 25 mg dose may be administered at bedtime, if necessary Maximum dose: 125 mg per day Use: For acute or long-term use in the relief of signs and symptoms of ankylosing spondylitis. Usual Adult Dose for Dysmenorrhea: Diclofenac potassium. find more information

Table 4 Side effects observed and percentage per gender.

Side Effects	Percentage per gender	
	Males	Females
Water retention	100%	100%
Acne	86%	98%
Gynaecomastia	36%	-
Aggression	100%	69%
Hypertension	68%	56%
Cardiovascular disease	8%	1%
Palpitations	62%	20%
Cardiac muscle hypertrophy	2%	-
Virilization	-	79%
MIgraines	78%	89%
Impotence	96%	-
Stomach pain	46%	79%
Jaundice	4%	1%
Muscle rupture	8%	1%
Prostate enlargement	78%	-
Hair loss	68%	5%
Stunted growth	24%	46%
Insomnia	98%	100%
Infertility	86%	100%
Anaphylactic reactions	12%	2%
Psychological disorders	100%	100%

Initial dose: 25 mg orally once a day, increased after one week to 50 mg orally once a day Maintenance dose: 50 to 200 mg orally once a day. Usual Adult Dose for Social Anxiety Disorder: Initial dose: 25 mg orally once a day, increased after one week to 50 mg orally once a day Maintenance dose: 50 to 200 mg orally once a day. 1-3.3 mg/kg/day orally or divided every 12 hours; not to exceed 3.3 mg/kg/day or up to 100 mg/day. Congestive Heart Failure. Adult and Geriatric. Indicated for NYHA class II/IV heart failure (provided CrCl greater than 30 mL/min and serum K less than 5 mEq/dL) Initial: 25 mg orally once/day.



Metoprolol Tartrate Immediate Release Tablets: Initial dose: 100 mg orally per day in single or divided doses Maintenance dose: 100 to 450 mg orally per day. Metoprolol Succinate Extended Release Tablets: Initial dose: 25 to 100 mg orally once a day Maintenance dose: 100 to 400 mg orally once a day. Usual Adult Dose for Angina Pectoris: Initial.



Maintenance dose: 0.125 to 0.25 mg orally once a day at bedtime Maximum dose: 0.25 mg/day Duration of therapy: 7 to 10 days Comments:-Failure to respond after 7 to 10 days may suggest a primary psychiatric and/or medical illness that requires further evaluation.-Use of this drug for more than 2 to 3 weeks requires complete reevaluation of the. visite site